

LECTURE #8

GENERAL MARKOV PROCESS IN EQUILIBRIUM

204528

Queueing Theory and
Applications in Networks

Assoc. Prof. Anan Phonphoem, Ph.D. (รศ.ดร. อนันต์ พลเพิ่ม)
Computer Engineering Department, Kasetsart University

Outline


- General Markov Process
- Erlangian Distribution
 - 2-stage Erlangian E_2
 - r-stage Erlangian E_r
- $M/E_r/1$

General Markov Process

- More general than birth-death process
 - Transition beyond nearest neighbors are allowed
 - But not too complicated
- We are interested in the Equilibrium Solutions

Simple three-state Markov Chain

4



- From Flow conservation
 - Prob. Flow in = Prob. Flow out

- For state 0:


- For state 1:

- For state 2:

$$(3/2)\lambda p_0 = \mu p_1$$

$$(\lambda + \mu)p_1 = \lambda p_0 + \mu p_2$$

$$\mu p_2 = (\lambda/2)p_0 + \lambda p_1$$



Simple three-state Markov Chain

- We need one more equation

$$p_0 + p_1 + p_2 = 1$$

- Solution

$$p_0 = [1 + 2(\lambda/\mu) + (3/2)(\lambda/\mu)^2]^{-1}$$

$$p_1 = 3/2(\lambda/\mu) p_0$$

$$p_2 = [(1/2)(\lambda/\mu) + (3/2)(\lambda/\mu)^2] p_0$$

Limiting Probability

- As $t \rightarrow \infty$
 - We can find the p_k
- How about prob. that an arriving customer finds the system in state E_k ?
 - It is equal to p_k , isn't it?

D/D/1

7

- Arrival is deterministic and occurs at every interval = \bar{t} sec
- Service time is also deterministic and identical for all customers = \bar{x} sec
- Let
 $r_k =$ [prob. that an arriving customer finds the system in state E_k]

D/D/1

8

- For stability of the system
 - It requires: $\bar{x} < \bar{t}$
 - $\rho = \bar{x} / \bar{t}$
- No customer has to wait !
 - $r_0 = 1$ and $r_k = 0$ for $k \geq 1$
- Therefore
 - $p_0 = 1 - \rho$ and $p_1 = \rho$
 - $p_k = 0$ for $k \geq 2$

D/D/1

- In summary
 - For D/D/1: $r_k \neq p_k$
- How about the other systems?
 - Many contains the property of $r_k = p_k$
 - Poisson arrival system is an example

Poisson Arrivals

- For

$P_k(t)$ = Prob. that system is in state E_k

$R_k(t)$ = Prob. that an arriving customer finds
the system is in state E_k

- Let $A(t, t+\Delta t)$ = Event that arrival occurs in the interval $(t, t+\Delta t)$

Poisson Arrivals

11

- $$\begin{aligned} R_k(t) &\stackrel{\Delta}{=} \lim_{\Delta t \rightarrow 0} P[N(t) = k \mid A(t, t+\Delta t)] \\ &= \lim_{\Delta t \rightarrow 0} \frac{P[N(t) = k, A(t, t+\Delta t)]}{P[A(t, t+\Delta t)]} \\ &= \lim_{\Delta t \rightarrow 0} \frac{P[A(t, t+\Delta t) \mid N(t) = k] P[N(t) = k]}{P[A(t, t+\Delta t)]} \\ &= \lim_{\Delta t \rightarrow 0} P[N(t) = k] \end{aligned}$$

$$P_k(t) = R_k(t)$$

Erlangian Distribution E_r

12

- A.K. Erlang proposed
 - The extreme simplicity of exponential distribution
 - Solving Markovian queueing systems
- The exponential distribution is not **always** appropriate for representing the true situation with regards to service times
- More general service distribution → Markov chain becomes complicated.




Agner Krarup Erlang
Denmark
From: Wikipedia

Erlangian Distribution Er

- Erlang purposed
 - The decomposing the service time distribution into a collection of exponential distributions

An exponential server

14



- For an exponentially distributed service time


$$\begin{aligned} b(x) &\triangleq \frac{dB(x)}{dx} \\ &= \mu e^{-\mu x} \quad x \geq 0 \end{aligned}$$

- For Mean and Variance of the service time

$$E[x] = \frac{1}{\mu} \quad \sigma_b^2 = \frac{1}{\mu^2}$$

A series of two exponential servers

15



- A series of two small exponential servers
- Each internal server has parameter = 2μ
- So the pdf of each server

$$h(y) = 2\mu e^{-2\mu y} \quad y \geq 0$$

A series of two exponential servers

- For Mean and Variance of the service time

$$E[x] = \frac{1}{2\mu} \quad \sigma_h^2 = \frac{1}{(2\mu)^2}$$

- The functions of the system
 - Upon departure of customer, a new customer is allowed to enter the service facility
 - After finish stage 1 → enter stage 2
 - Only one customer is allowed in the service facility

A series of two exponential servers

- What is the distribution of the time spent in the service facility ?
 - Sum of the two independent random variables
 - The convolution of the density function of the two random variables

Convolution of two RVs

- $Y = X_1 + X_2$
- $F_Y(y) = P[Y \leq y] = P[X_1 + X_2 \leq y]$
$$= \int_{-\infty}^{\infty} \int_{-\infty}^{y-x_2} f_{X_1, X_2}(x_1, x_2) dx_1 dx_2$$
$$= \int_{-\infty}^{\infty} \left[\int_{-\infty}^{y-x_2} f_{X_1}(x_1) dx_1 \right] f_{X_2}(x_2) dx_2$$
$$= \int_{-\infty}^{\infty} F_{X_1}(y - x_2) f_{X_2}(x_2) dx_2$$


Convolution of two RVs

- $F_Y(y) = \int_{-\infty}^{\infty} F_{X_1}(y - x_2) f_{X_2}(x_2) dx_2$
- $f_Y(y) = \int_{-\infty}^{\infty} f_{X_1}(y - x_2) f_{X_2}(x_2) dx_2$
- $f_Y(y) = f_{X_1}(y) \otimes f_{X_2}(y)$

Convolution of two RVs

$$(f * g)(t) \stackrel{\text{def}}{=} \int_{-\infty}^{\infty} f(\tau) g(t - \tau) d\tau$$

- Express each function in terms of a dummy variable τ .
- Reflect one of the functions: $g(\tau) \rightarrow g(-\tau)$.
- Add a time-offset, t , which allows $g(t - \tau)$ to slide along the τ -axis.
- Start t at $-\infty$ and slide it all the way to $+\infty$. Wherever the two functions intersect, find the integral of their product.




Laplace transform in probability

- In probability, the Laplace transform is defined by means of an expectation value.
- If X is a random variable with probability density function f ,
 - then the Laplace transform of f is given by the expectation

$$\begin{aligned}(\mathcal{L}f)(s) &= E[e^{-sX}] . \\ &= \int_0^{\infty} e^{-sx} f_X(x) dx\end{aligned}$$

Laplace transform of pdf

22



Service Facility

- $B^*(s) \triangleq \int_0^{\infty} e^{-sx} b(x) dx$
- $H^*(s) \triangleq \int_0^{\infty} e^{-sy} h(y) dy$
- $B^*(s) = H^*(s) H^*(s) = [H^*(s)]^2$

2-stage Erlangian Server E_2

23

- $h(y) = 2\mu e^{-2\mu y}$
- $H^*(s) \triangleq \int_0^{\infty} e^{-sy} h(y) dy$
 $= \frac{2\mu}{s + 2\mu}$

- $B^*(s) = [H^*(s)]^2$
 $= \left(\frac{2\mu}{s + 2\mu} \right)^2$


Inverse

$$b(x) = 2\mu(2\mu x) e^{-2\mu x}$$

For $x \geq 0$

A series of two exponential servers

24



- For Mean and Variance of the service time

$$E[x] = 2E[y] = \frac{1}{\mu}$$


$$\sigma_b^2 = \sigma_h^2 + \sigma_h^2 = \frac{1}{2\mu^2}$$

A series of two exponential servers

- Mean time for single and two-stage are the same
 - By speed up the internal service 2 times
- Variance of the two-stage is one-half of the single server

r-stage Erlangian server E_r

26



Service Facility

- $h(y) = r\mu e^{-r\mu y} \quad y \geq 0$
- For Mean and Variance of each stage

$$E[y] = \frac{1}{r\mu}$$
$$\sigma_h^2 = \left(\frac{1}{r\mu} \right)^2$$

r-stage Erlangian server Er

- For Mean and Variance of the service time

$$E[x] = rE[y] = \frac{1}{\mu}$$

$$\sigma_b^2 = r \left(\frac{1}{r\mu} \right)^2 = \frac{1}{r\mu^2}$$

r-stage Erlangian server E_r

28

- Solve for the pdf of the service time


$$B^*(s) = \left(\frac{r\mu}{s + r\mu} \right)^r$$

$$b(x) = \frac{r\mu(r\mu x)^{r-1} e^{-r\mu x}}{(r-1)!} \quad x \geq 0$$

Erlang Distribution

r-stage Erlangian distribution E_r

29



r-stage Erlangian distribution E_r

- Note
 - $r \uparrow \rightarrow$ pdf approaches Gaussian Distribution
 - Mean is constant
 - The standard deviation (width) shrinks by $1/\sqrt{r}$
 - If $r \rightarrow \infty$ then
pdf \rightarrow unit impulse function at $x = 1/\mu$

M / E_r / 1

31

- Memoryless / Erlangian/1 server
- Define

$$a(t) = \lambda e^{-\lambda t} \quad t \geq 0$$

$$b(x) = \frac{r\mu(r\mu x)^{r-1} e^{-r\mu x}}{(r-1)!} \quad x \geq 0$$


M / E_r / 1

32

- State variable = total number of service stage yet to be completed
- k = # of customers
- Now we are in the i^{th} stage of service ($1, r$)
- j = number of stages left in the total system
 - = $(k - 1)r + (r - i + 1)$
 - = $rk - i + 1$

of stages left to be completed

33




- $j = rk - i + 1$
- For $r = 3$ stages, 1 customer, now in stage 2
$$j = 3 * 1 - 2 + 1 = 2$$

= “there are **2 stages** left to be completed”

of stages left to be completed


34



- $j = rk - i + 1$
- For $r = 3$ stages, 3 customers, now in stage 1
 $j = 3*3 - 1 + 1 = 9$
= “there are **9 stages** left to be completed”

of stages left to be completed

35



- $j = rk - i + 1$
- For $r = 3$ stages, 3 customers, now in stage 3
$$j = 3 * 3 - 3 + 1 = 7$$

= “there are **7 stages** left to be completed”


M / E_r / 1

36

- p_k = equilibrium prob. for the # of customers in the system
- $P_j = P[j \text{ stages in the system}]$
- $p_k = \sum_{j=(k-1)r+1}^{kr} P_j \quad k = 1, 2, 3, \dots$

M / E_r / 1 State diagram

37



- $P_j = 0 \quad j < 0$
- $\lambda P_0 = r\mu P_1$
- $(\lambda + r\mu)P_j = \lambda P_{j-r} + r\mu P_{j+1} \quad j = 1, 2, \dots$

M / E_r / 1

38

- Solved by using z -transform

$$P(z) = \sum_{j=0}^{\infty} P_j z^j$$

- Multiplied by z^j and sum

$$\sum_{j=1}^{\infty} (\lambda + r\mu) P_j z^j = \sum_{j=1}^{\infty} \lambda P_{j-r} z^j + \sum_{j=1}^{\infty} r\mu P_{j+1} z^j$$

$$(\lambda + r\mu) \left(\sum_{j=0}^{\infty} P_j z^j - P_0 \right) = \lambda z^r \sum_{j=1}^{\infty} P_{j-r} z^{j-r} + (r\mu/z) \sum_{j=1}^{\infty} P_{j+1} z^{j+1}$$

M / E_r / 1

39

$$(\lambda + r\mu) [P(z) - P_0] = \lambda z^r P(z) + (r\mu/z) [P(z) - P_0 - P_1 z]$$

$$P(z) = \frac{r\mu P_0(1 - z)}{r\mu + \lambda z^{r+1} - (\lambda + r\mu)z}$$

Finding P_0

$$P(1) = 1 = \frac{r\mu P_0}{r\mu - \lambda r}$$

$$P_0 = 1 - (\lambda/\mu) \rightarrow \rho$$

M / E_r / 1

40

$$P(z) = \frac{r\mu(1 - \rho)(1 - z)}{r\mu + \lambda z^{r+1} - (\lambda + r\mu)z}$$

- Then invert the z -transform
→ Distribution of # of the stages in the system

$M / E_r / 1$ (For $r = 1$)

41

- For case $r = 1$

$$\begin{aligned} P(z) &= \frac{\mu(1 - \rho)(1 - z)}{\mu + \lambda z^2 - (\lambda + \mu)z} \\ &= \frac{(1 - \rho)(1 - z)}{1 + \rho z^2 - (1 + \rho)z} \\ &= \frac{(1 - \rho)}{(1 - \rho z)} \end{aligned}$$

M / E_r / 1 (For r = 1)

42

- Invert z-transform

$$P_k = (1 - \rho) \rho^k \quad k = 0, 1, 2, \dots$$

- $$p_k = \sum_{j=(k-1)r+1}^{kr} P_j \quad k = 1, 2, 3, \dots$$

- For r = 1 $p_k = P_k \rightarrow M / M / 1$

M / E_r / 1 (For general r)

43

$$\begin{aligned} P(z) &= \frac{r\mu(1-\rho)(1-z)}{r\mu + \lambda z^{r+1} - (\lambda + r\mu)z} \\ &= \frac{(1-\rho)}{(1-z/z_1)(1-z/z_2)\dots(1-z/z_r)} \\ &= (1-\rho) \sum_{i=1}^r \frac{A_i}{(1-z/z_i)} \\ A_i &= \prod_{\substack{n=1 \\ n \neq i}}^r \frac{1}{(1-z_i/z_n)} \end{aligned}$$

M / E_r / 1 (For general r)

44

- Invert z-transform

$$P_j = (1 - \rho) \sum_{i=1}^r A_i (z_i)^{-j} \quad j = 1, 2, \dots, r$$

$E_r / M / 1$

45


- Erlangian / Memoryless / 1 server
- Define

$$a(t) = \frac{r\lambda(r\lambda t)^{r-1}e^{-r\lambda t}}{(r-1)!} \quad t \geq 0$$

$$b(x) = \mu e^{-\mu x} \quad x \geq 0$$

$E_r / M / 1$

46



$$\text{Arriving : } a(t) = \frac{r\lambda(r\lambda t)^{r-1}e^{-r\lambda t}}{(r-1)!} \quad t \geq 0$$

$$\text{Service : } b(x) = \mu e^{-\mu x} \quad x \geq 0$$

$E_r / M / 1$

47


- State variable = total number of service stage yet to be completed
- k = # of customers
- Now the arriving customer is in the i^{th} stage of service $[1, r]$
- j = total number of stages of arrival in the system
= $rk + i - 1$

$E_r / M / 1$

- p_k = equilibrium prob. for the # of customers in the system
- P_j = # of arrival stages in the system
- $p_k = \sum_{j=rk}^{r(k+1)-1} P_j$

$E_r / M / 1$ State diagram

49



- Each departure = remove “ r ” stages of arrival
- $r\lambda P_0 = \mu P_r$
- $r\lambda P_0 = r\lambda P_{j-1} + \mu P_{j+r} \quad 1 \leq j \leq r-1$
- $(r\lambda + \mu)P_j = r\lambda P_{j-1} + \mu P_{j+r} \quad r \leq j$

$E_r / M / 1$

50

- Solved by using z -transform

$$P(z) = \sum_{j=0}^{\infty} P_j z^j$$

- Multiplied by z^j and sum
- For $j \geq 1$

$$\sum_{j=1}^{\infty} (\mu + r\lambda) P_j z^j - \sum_{j=1}^{r-1} \mu P_j z^j = \sum_{j=1}^{\infty} r\lambda P_{j-1} z^j + \sum_{j=1}^{\infty} \mu P_{j+r} z^j$$

$E_r / M / 1$

51

$$P(z) = \frac{(1 - z^r) \sum_{j=1}^{r-1} P_j z^j}{r \rho z^{r+1} - (1 + r\rho)z^r + 1}$$

$$P(z) = \frac{(1 - z^r)(1 - 1/z_0)}{r(1 - z)(1 - z/z_0)}$$

$$P_j = \begin{cases} \frac{1}{r}(1 - z_0^{-j-1}) & 0 \leq j < r \\ \rho(z_0 - 1)z_0^{r-j-1} & j \geq r \end{cases}$$

$E_r / M / 1$

52

$$p_k = \begin{cases} 1 - \rho & k = 0 \\ \rho(z_0^r - 1)z_0^{-rk} & k > 0 \end{cases}$$